

- Nurse, C. G., 1910b. Further notes regarding the breeding of *Chilosia grossa*. *Entomologist*. **43**: 313-314.
- Réaumur, R. A. F. de., 1740. *Mémoires pour servir à l'histoire des Insectes. V. Paris*.
- Roser, C. L. F. von., 1834. Verzeichnisse in Wörtemberg vorkommender zweiflüglicher Insekten. *KorrespBl. k. Wurrt. Landwirt ver Stuttgart*. **1834**: 1-19.
- Rossi, F., 1848. *Systematisches verzeichniss der Zweiflügelichten Insecten (Dipiera) des Erzherzogthumes Österreich*. Wien.
- Trägårdh, J., 1939. *Sveriges Skoginsepter 2*. Auflage. Stockholm.
- Verrall, G. H., 1901. *British Flies. 8. Platypezidae, Pipunculidae and Syrphidae of Great Britain*. London.
- Vimmer, A., 1925. *Larvy a kukly dvojkrídleho Stredoevropského se zvláštěm z retelem na skudce rostlin kulturnich*. Prague.
- Weyenberg, H., 1869. Nederlandsche Diptera in Metamorphose en levenswijze. *Tijds. v. Ent.* **12**: 155-174.
- Zetterstedt, J. W., 1843. *Diptera Scandinaviae II*. Lund.

Postscript

Since writing the above paper Mr. Alan Stubbs has kindly drawn my attention to two records of reared *Cheilosia semifasciata* Becker: Speight, M. C. D., Chandler, P. J. & Nash, R., 1975, *Proc. R. Irish. Acad.* **75** (B) (1): 9, record it mining *Umbilicus*; Uffen, R. & Chandler, P. J., 1978, *In Subbs, A. E. & Chandler, P. J., A Dipterist's Handbook, Amat. Ent.* **15**: 221, record it from *Sedum telephium* and *Umbilicus*.

GEGENES PUMILIO (LEP.: HESPERIIDAE): A RECORD FOR CRETE. — Belatedly, I would like to record the capture of a single specimen of this species on the island of Crete. The specimen was taken on 9.iv.1973 on rough land, close to a small patch of cultivated land at about 400 metres above sea-level, close to Neapolis in the east of the island. This area is about 20 kilometres north of the Plateau of Lasithi, and 10 kilometres from the north coast of the island. Whilst this is not the first record of this butterfly occurring in Crete (L. G. Higgins *pers. comm.*), the fact remains unrecorded in Higgins and Riley, *A Field Guide to the Butterflies of Britain and Europe*, third Edition (1975). — DAVID C. HOCKIN, Culterty Field Station, University of Aberdeen, Newburgh, Ellon, Aberdeenshire.

KYBOASCA BIPUNCTATA (OSHANIN) (HOMOPTERA: AUCHEGORHYNCHA: TYPHLOCYBINAЕ): A SPECIES NEW TO BRITAIN. — 40 ♂ 40 ♀ specimens of *Kyboasca bipunctata* (Oshanin) were identified in a large sample of leafhoppers collected from English Elm (*Ulmus procera*) at Mitcham, Surrey on 2nd July 1978. Although previously unrecorded in Britain, *K. bipunctata* is widely recorded in the Palaearctic on Elms. A further male specimen was taken on English Elm at Twitton nr. Otford, Kent, July 1978, by Mr. W. R. Dolling. My thanks to Dr. J. Dlabola (Narodni Museum, Czechoslovakia) for confirming the identification. It will be dealt with by Dr. W. J. de Le Quesne in his forthcoming Royal Entomological Society handbook on the Typhlocybinae. — M. R. WILSON, Department of Zoology, University College, Cardiff.